

FULANO DE TAL

Num. Atend. :

Idade em:

Médico Dr(a): Conv.:

Data :

Dt. Nasc.:

Sexo:

Entrega.:

HORMÔNIO SALIVA – PERFIL AVANÇADO PLUS

Report Number:

Provider:

Ordering Provider:


Patient Info:

Age: Gender:

Menopausal Status:

Sample Collection Date/Time

	Saliva Hormone Test	Result	Units	L	WR	H	Reference Range
HORMONES	Estrone (E1)	6.00	pg/ml				7.2-40.6 pre menopausal
	Estradiol (E2)	6.48	pg/ml		◆		1.5-10.8 supplementation
	Estriol (E3)	30.00	pg/ml		◆		<30.0 female, non-pregnant
	EQ (E3 / (E1 + E2))						
	Progesterone (Pg)	200.00	pg/ml				127.0-446.0 pre menopausal (luteal)
	Ratio of Pg/E2						
ADRENALS	Testosterone	56.00	pg/ml			↑	6.1-49.0 female
	DHEA	252.60	pg/ml		◆		106.0-300.0 female
	Cortisol Morning	41.00	nmol/L			↑	5.1-40.2; optimal range: 18-35*
	Cortisol Noon	15.00	nmol/L		◆		2.1-15.7; optimal range: 6-12*
	Cortisol Evening	2.00	nmol/L		◆		1.8-12; optimal range: 4-8*
	Cortisol Night	0.80	nmol/L	↓			0.9-9.2; optimal range: 2-6*



Hormone Interpretations:

- Estrone and estradiol are within the reference ranges, however the Estrogen Quotient (EQ) is suboptimal. Estriol is less potent than the other estrogens and when present in sufficient quantities (as indicated by an optimal EQ) it plays an antagonistic role, and may govern the proliferative effects of estrone and estradiol. Although estriol level is above the reference range (likely do to individual variance), estriol supplementation is a consideration to optimize this quotient and reduce associated risks. * References available upon request.
- Progesterone to estradiol (Pg/E2) ratio and reported symptoms are consistent with estrogen dominance. Supplementation with topical progesterone to correct this relative deficiency is a consideration.
- DHEA level is consistent with stress response or supplementation (not reported), although metabolic syndrome cannot be ruled out. Serum vitamin D, fasting glucose and insulin testing may be warranted.
- Adrenal gland function appears reasonably adequate. Query thyroid insufficiency (perhaps related to iodine deficiency).

Notes:

L=Low(below range) WR=Within Range (within range) H=High (above range)

DHEA, Testosterone, Estrone and Estriol results are for investigational use only.

*Apply only when all four cortisol are measured. Clinical interpretations may override these generalized optimal ref. ranges.

**The Pg/E2 ratio is an optimal range established based on clinical observation. Progesterone supplementation is generally required to achieve this level in men and postmenopausal women.

Adrenal Phase:

